

Generalized Quantum Search Hamiltonian

Joonwoo Bae*, Younghun Kwon†

Department of Physics, Hanyang University,
Ansan, Kyunggi-Do, 425-791, South Korea

February 1, 2008

Abstract

There are hamiltonians that solve a search problem of finding one of N items in $O(\sqrt{N})$ steps. They are hamiltonians to describe an oscillation between two states. In this paper we propose a generalized search hamiltonian, H_g . Then the known search hamiltonians become special cases of H_g . From the generalized search hamiltonian, we present remarkable results that searching with 100% is subject only to the phase factor in H_g and independent to the number of states or initialization.

Grover proposed the quantum search algorithm that finds one of unsorted N items in $O(\sqrt{N})$ steps.[1] Since it is known that classical search algorithms would take $O(N)$ steps to solve a search problem, the quadratic speedup in Grover algorithm is remarkable. The reason of the speedup is effects of quantum mechanics. To apply quantum mechanics to a search problem, it needs to map each item to a state, respectively, in N dimensional Hilbert space. Then a search problem can be transformed to finding one of N states. We can use quantum mechanical characteristics, for instance, superposition and parallelism. Grover algorithm is in fact to find the state which corresponds to a target item. An iteration of Grover algorithm is composed of two operations: the first is to flip the target state about 0 and the next to invert all states about the average state.[2] An oracle, the heart of quadratic speedup, is applied in the first operation. Grover iterations amplify the amplitude of the target state up to nearly one in $O(\sqrt{N})$ times, when an initial state is a uniform superposition of N states. Zalka showed Grover algorithm is optimal when unstructured items are considered.[3]

On the other hand, there are a search algorithm based on hamiltonian evolution by Schrodinger equation. While Grover algorithm operates a state in discrete time, a search hamiltonian does a state in continuous time. They are hamiltonians to describe an oscillation between two states. Farhi et al. suggested harmonic-oscillation hamiltonian, exactly $H_{fa} = E(|w\rangle\langle w| + |s\rangle\langle s|)$, where $|w\rangle$ is a target state and $|s\rangle$ is an initial state.[4] Fenner provided another hamiltonian, $H_{fe} = 2iEx(|w\rangle\langle\psi| - |\psi\rangle\langle w|)$, where x is the overlap between an initial state and a target state, i.e., $\langle w|s\rangle = x(> 0)$. [5] A state under the hamiltonians evolves from an initial state to a target state in the half period of oscillation,

*Email address: jwbae@newton.hanyang.ac.kr

†Email address: yhkwon@newton.hanyang.ac.kr

$O(\sqrt{N})$. This oscillation is a process to amplify the amplitude of a target state. It is noted that the initial state should be prepared as a superposition of all states, since one of initially superposed states becomes a target state by the hamiltonian evolution.

We expect that there exists a more general search hamiltonian including the known hamiltonians. We then write down a most possible combination of a target state and an initial state as follows:

$$H = E(a|w\rangle\langle w| + b|w\rangle\langle s| + c|s\rangle\langle w| + d|s\rangle\langle s|)$$

The initial state can be written as $|s\rangle = x|w\rangle + \sqrt{1-x^2}|r\rangle$, where $|r\rangle$ is the state orthogonal to the target state (so $\langle w|r\rangle = 0$) and x is the overlap between the initial state and the target state, i.e., $x = \langle w|s\rangle$. The hamiltonian and the initial state can be written in matrix representation:

$$\mathbf{H} = E \begin{pmatrix} a + (b+c)x + dx^2 & (b+dx)\sqrt{1-x^2} \\ (c+dx)\sqrt{1-x^2} & d(1-x^2) \end{pmatrix},$$

$$|\mathbf{s}\rangle = \begin{pmatrix} x \\ \sqrt{1-x^2} \end{pmatrix}$$

A state $|\psi(t)\rangle$ is governed by time-dependent Schrodinger equation:

$$i\frac{d}{dt}|\psi(t)\rangle = H|\psi(t)\rangle$$

where the initial condition is $|\psi(t=0)\rangle = |s\rangle$ (we let $\hbar = 1$ for convenience). After t times the probability to find the target state is :

$$\begin{aligned} P &= |\langle w|e^{-iHt}|s\rangle|^2 \\ &= x^2 \cos^2 EDT + CC^* \sin^2 EDT \end{aligned} \quad (1)$$

where, $q = \frac{1}{2}((a+d) + (b+c)x)$,
 $D = \sqrt{q^2 - (ad-bc)(1-x^2)}$, and $C =$

$$\left[\frac{(d(1-x^2) - q)^2 + x(c+dx)(d(1-x^2) - q) - D^2}{(c+dx)D} \right]$$

The postulate of quantum mechanics, that any observable should be hermitian, requires $H = H^\dagger$. Then we obtain :

1. a and d are real.
2. $b = c^* = re^{i\phi}$, where r and ϕ are real.

Suppose the initial state is a uniform superposition of all states and we read out the final state at time T . In equation (1), the contribution of $x^2 \cos^2 EDT$ to the probability is so small that it can be negligible. Moreover, $x^2 \cos^2 EDT$ becomes zero at time T , since the read-out time T comes from $EDT = \frac{\pi}{2}$. We find out $T = O(\sqrt{N})$, where the quadratic speedup is shown. Thus it is essential

to make the value of C^*C as large as possible, to find the target state with high probability at time T .

$$P = CC^* = \frac{u_0 + u_1x + u_2x^2 + u_3x^3 + u_4x^4 + u_5x^5 + u_6x^6}{l_0 + l_1x + l_2x^2 + l_3x^3 + l_4x^4} \quad (2)$$

$$\begin{aligned} u_0 &= r^4 \\ u_1 &= (a + 3d)r^3 \cos \phi \\ u_2 &= \frac{1}{4}r^2(a^2 + 6ad + 9d^2 - 4r^2 \\ &\quad + 4(ad + d^2 + r^2)\cos 2\phi) \\ u_3 &= \frac{1}{2}dr \cos \phi(a^2 + 4ad + 3d^2 - 4r^2 + 4r^2 \cos 2\phi) \\ u_4 &= \frac{1}{4}(a^2d^2 + 2ad^3 + d^4 - 4d^2r^2 + 2r^4 \\ &\quad + (4d^2r^2 - 2r^4)\cos 2\phi) \\ u_5 &= 2dr^3 \cos \phi \sin^2 \phi \\ u_6 &= d^2r^2 \sin^2 \phi \\ l_0 &= \frac{1}{4}(a^2r^2 - 2adr^2 + d^2r^2 + 4r^4) \\ l_1 &= \frac{1}{2}r(a^2d - 2ad^2 + d^3 + 2ar^2 + 6dr^2)\cos \phi \\ l_2 &= \frac{1}{4}(a^2d^2 + d^4 + 8d^2r^2 - 2r^4 - 2a(d^3 - 4dr^2) \\ &\quad + 2r^2(2ad + 2d^2 + r^2)\cos 2\phi) \\ l_3 &= dr \cos \phi(3ad + d^2 - r^2 + r^2 \cos 2\phi) \\ l_4 &= \frac{1}{2}d^2(2ad - r^2 + r^2 \cos 2\phi) \end{aligned} \quad (3)$$

In (2), to consider the zeroth order terms of x , u_0 and l_0 , is crucial. Thus we obtain a condition $a = d$ by setting $u_0 = l_0$, which automatically gives $u_1 = l_1$. Under the condition $a = d$, the probability becomes:

$$P_g = 1 - O(x^2) \quad (4)$$

The corresponding hamiltonian is:

$$H_g = E_1(|w\rangle\langle w| + |s\rangle\langle s|) + E_2(e^{i\phi}|w\rangle\langle s| + e^{-i\phi}|s\rangle\langle w|) \quad (5)$$

where $E_1 = Ea$ and $E_2 = Er$. The probability to get the target state under this hamiltonian is not exactly but nearly one. Therefore it is allowed to use hamiltonian H_g when searching with 100% is not required.

We here pursue condition for higher probability, which is $l_2 = u_2$. It makes us choose a specific phase, $\phi = n\pi$. This automatically gives useful relations, $l_3 = u_3, l_4 = u_4$, and $u_5 = u_6 = 0$. Then the probability under these relations is:

$$P_p = \frac{u_0 + u_1x + u_2x^2 + u_3x^3 + u_4x^4}{u_0 + u_1x + u_2x^2 + u_3x^3 + u_4x^4} = 1$$

The hamiltonian is :

$$H_p = E_1(|w\rangle\langle w| + |s\rangle\langle s|) \pm E_2(|w\rangle\langle s| + |s\rangle\langle w|) \quad (6)$$

That is, hamiltonian H_p can search a target state with probability one.

We now have the generalized search hamiltonian, H_g , and the perfect search hamiltonian H_p . H_p is in particular produced from H_g , by fixing the phase $\phi = n\pi$. Here, we can show that known search hamiltonians are special forms of H_g . For $E_2 = 0$, it becomes Farhi's hamiltonian, and Fenner's hamiltonian for $E_1 = 0$, $\phi = \frac{\pi}{2}$, and $r = 2x$. Moreover, we provide a new quantum search hamiltonian :

$$H_{new} = E_2(e^{i\phi}|w\rangle\langle\psi| + e^{-i\phi}|\psi\rangle\langle w|) \quad (7)$$

When Grover algorithm is applied to a search problem, the probability one appears if and only if two assumptions are satisfied : 1) the target items are $\frac{N}{4}$ among N items, 2) the initial state is a uniform superposition of all states in a system. Here we present the remarkable result that hamiltonian H_p does not depend on the number of states and initialization. There is no condition on H_p about the number of states or initialization. This implies that H_p finds target states with probability one regardless of the number of target states, though the initial state is prepared as an arbitrary superposition of all states. When a generalized search hamiltonian is used in a search problem, probability one at read-out procedure depends only on the phase condition.

Acknowledgement

J. Bae is supported in part by the Hanyang University Fellowship and Y. Kwon is supported in part by the Fund of Hanyang University.

References

- [1] L. K. Grover, "Quantum Mechanics helps in searching for a needle in a haystack", *Phys. Rev. Letters*, 78(2), 325, 1997.
- [2] Richard Jozsa, "Searching in Grover's Algorithm", <http://xxx.lanl.gov/abs/quant-ph/9901021>.
- [3] C. Zalka, "Grover's quantum searching algorithm is optimal", *Phys. Rev. A* 60(1999), 2746-2751, <http://xxx.lanl.gov/abs/quant-ph/9711070>.

- [4] E. Farhi and S. Gutmann, “An Analog Analogue of a Digital Quantum Computation”, Phys. Rev. A 57, 2403 (1998), <http://xxx.lanl.gov/abs/quant-ph/96120026>.
- [5] Stephen A. Fenner, “An intuitive Hamiltonian for quantum search”, <http://xxx.lanl.gov/abs/quant-ph/0004091>.